REVIEW ARTICLE

Check for updates

WILEY

Biochar affects greenhouse gas emissions in various environments: A critical review



¹Tianjin Key Laboratory of Clean Energy and Pollution Control, School of Energy and Environmental Engineering, Hebei University of Technology, Tianjin, PR China

²Key Laboratory of Pollution Processes and Environmental Criteria (Ministry of Education), Tianjin Engineering Research Center of Environmental Diagnosis and Contamination Remediation, College of Environmental Science and Engineering, Nankai University, Tianjin, PR China

³Department of Renewable Resources, University of Alberta, Edmonton, Alberta, Canada

⁴UK Biochar Research Centre, School of GeoSciences, University of Edinburgh, Edinburgh, UK

⁵Korea Biochar Research Center & APRU Sustainable Waste Management Program & Division of Environmental Science and Ecological Engineering, Korea University, Seoul, South Korea

Correspondence

Jingchun Tang, College of Environmental Science and Engineering, Nankai University, Tianjin 300350, PR China. Email: tangjch@nankai.edu.cn

Yong Sik Ok, Korea University, Seoul 02841, South Korea. Email: yongsikok@korea.ac.kr

Funding information

National Natural Science Foundation of China, Grant/Award Numbers: 41877372, 42177218, U18062162; National Key R&D Program of China, Grant/Award Number: 2018YFC1802002; Hebei Outstanding Youth Science Foundation, Grant/Award Number: D2019202453; China Postdoctoral Science Foundation, Grant/Award Number: 2019M660967; the 111 Program, Ministry of Education, China, Grant/Award Number: T2017002: National Research Foundation of Korea (NRF) grant funded by the Korean Government (MSIT), Grant/Award Number: 2021R1A2C2011734; Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Grant/Award Number: NRF-2021R1A6A1A10045235: OJEong Resilience Institute, Korea University, Republic of Korea

Abstract

Biochar application to the soil is a novel approach to carbon sequestration. Biochar application affects the emission of greenhouse gases (GHGs), such as CO2, CH4, and N₂O, from different environments (e.g., upland soils, rice paddies and wetlands, and composting environments). In this review, the effect of biochar on GHGs emissions from the above three typical environments are critically evaluated based on a literature analysis. First, the properties of biochar and engineered biochar related to GHGs emissions was reviewed, targeting its relationship with climate change mitigation. Then, a meta-analysis was conducted to assess the effect of biochar on the emissions of CO₂, CH₄, and N₂O in different environments, and the relevant mechanisms. Several parameters were identified as the main influencing factors in the meta-analysis, including the pH of the biochar, feedstock type, pyrolysis temperature, biochar application rate, C/N ratio of the biochar, and experimental scale. An overall suppression effect among different environments was found, in the following order for different greenhouse gases: $N_2O > CH_4 > CO_2$. We conclude that biochar can change the physicochemical properties of soil and compost in different environments, which further shapes the microbial community in a specific environment. Biochar addition affects CO₂ emissions by influencing oligotrophic and copiotrophic bacteria; CH₄ emissions by regulating the abundance of functional genes, such as mcrA (a methanogen) and pmoA (a methanotroph); and N₂O emissions by controlling Ncycling functional genes, including amoA, nirS, nirK, nosZ. Finally, future research directions for mitigating greenhouse gas emissions through biochar application are suggested.

KEYWORDS

biochar, black carbon, gasification, pyrolysis, UN Sustainable Development Goals

1 | INTRODUCTION

Climate change has accelerated with industrial development and the need to address this challenge is widely accepted by society and policymakers. Several pathways to zero carbon (C), or even a negative-C future, have been charted; however, achieving these goals is an enormous task, requiring multilateral efforts and different approaches, including emission reductions, CO_2 capture, and atmospheric greenhouse gas (GHG) removal. Among the six major GHGs listed in the Climate Change Control Inventory, CO_2 , CH_4 , and N_2O contribute the most to global climate change, with relative contributions of 60%, 20%, and 10% respectively (Josep et al., 2019). The concentration of CO_2 in the atmosphere has increased from 280 ppm in the AD 1700s to over 400 ppm, reflecting a rapid increase in CO_2 emissions since the Industrial Revolution (Sriphirom et al., 2020). Various approaches aimed at mitigating or minimizing climate change have been proposed to address the rising emissions of GHGs and their concentrations in the atmosphere (Song et al., 2019).

Carbon sequestration can directly decrease the emission of CO₂ into the atmosphere, and a new class of technologies, GHG removal technologies, have emerged to aid in reducing GHG concentrations in the atmosphere. Biochar is one piece of this puzzle as it has considerable global potential to sequester atmospheric C. The ability of biochar to sequester C from the atmosphere by plants has been the driving force behind its development. Biochar production itself can offset GHG emissions because it converts the organic C in the feedstock into stable C to prevent the degradation of biomass from releasing CO2 and CH4 into the atmosphere (Zhang & Ok, 2014). The application of biochar is supposed to be able to offset a maximum of 12% of current anthropogenic CO₂-C equivalent (CO2-Ce) emissions (i.e., 1.8 Pg CO2-Ce per year of the 15.4 Pg CO₂-C_e emitted annually; 1 Pg = 1 Gt) (Woolf et al., 2010). As an important indicator of the effectiveness of C sequestration, the stability of biochar in different soils has been extensively studied (Lian & Xing, 2017), and it is now widely accepted that the stability of most of the C contained in biochar is of the order of hundreds or even thousands of years (Spokas & Reicosky, 2009).

Biochar is produced from different feedstocks and is widely used in various environmental processes. The main functions of biochar can be summarized as follows: (1) The production of biochar, combined with energy recovery, is a good method for managing agricultural waste and has been practised in China and around the world (Lee et al., 2017). (2) Biochar is widely used as a soil conditioner to improve soil quality and crop yield (Pariyar et al., 2020) because its porous structure can improve soil quality by enhancing soil aeration, reducing soil hardening, and increasing soil cation exchange capacity (CEC). In addition, the nutrient content of biochar is important for plant growth and crop yields. (3) Biochar can be used for the remediation of soil and water contaminants (Xiao et al., 2020). In addition, engineered biochars have been developed to enhance biochar functions, such as adsorption, reduction, oxidation, and catalysation of specific pollutants (Lyu et al., 2020). Biochar has also been applied to increase the efficiency of waste treatment processes such as composting. (4) Biochar can be used for C sequestration and as an adsorbent for GHGs, such as CO2, to mitigate climate change (Huang et al., 2015).

Biochar also plays an important role in mitigating climate change by regulating GHG emissions from the soil and different environmental processes. Biochar application can change soil properties and hence affect microbial biomass, community structure, and activity, resulting in changes in soil GHG emissions. As microbial communities in uplands are quite different from those in paddy soils and wetlands, the application of biochar will have a different effect on GHG emissions in these two environments. For example, CH₄ emissions from rice paddy fields are much higher than those from upland fields, and the emissions from a rice paddy in the monsoon season in Asia account for $\sim 25\%$ -36% of global CH₄ emissions (Zhang, Qu, et al., 2020) because of extensive rice cultivation. It has been estimated that the application of biochar to paddy soils reduces seasonal CH₄ emissions by 40% (Sriphirom et al., 2020). Emissions of CO₂ are the main concern in upland agriculture, where biochar can reduce the net ecosystem CO2 exchange in crop production by 144%-283% (Azeem et al., 2019). Another GHG is N₂O, which has a much higher global warming potential and can be a key factor in both paddy and upland fields (Aamer et al., 2020). Although biochar generally reduces N₂O emissions from soil (Thangarajan et al., 2018), in some cases, it can enhance N₂O emissions from upland fields when water content increases (Troy et al., 2013). In addition to paddy and upland fields, biochar may also affect GHG emissions from industrial sites such as composting, anaerobic digestion, and bioremediation sites.

To date, several review papers have been published that focus on the effects of biochar on soil GHG emissions. These studies have summarized the effect of biochar on the properties and GHG emissions in the soils of a certain type of environment, such as forest soils (Li et al., 2018) or agricultural soils (Sri et al., 2021). However, no systematic review has compared the effects of biochar on GHG emissions from microbial processes in various environments (e.g., upland soils, rice paddies and wetlands, and composting environments), which is important for mitigating GHG emissions and promoting the application of biochar. In this study, we systematically evaluated the effects of biochar on GHG emissions in various environments (i.e., upland soils, rice paddies and wetlands, and composting environments) and the mechanisms involved. First, recent research and development on biochar production related to climate change mitigation are summarized. Second, the effects of biochar application on GHG emissions in upland fields, rice paddies and wetlands, compost systems, and the mechanisms involved (including the mechanisms that control GHG emissions based on the effects of biochar on soil physicochemical and microbial properties) are summarized.

2 | PROPERTIES OF PRISTINE AND ENGINEERED BIOCHAR RELEVANT TO CLIMATE CHANGE MITIGATION

2.1 | Properties of biochar relevant to climate change mitigation

Biochars have been widely applied to soil improvement in various environments, including uplands, rice paddies, wetlands, and

099145x, 2022, 17, Downloaded from https:

//onlinelibrary.wiley.com/doi/10.1002/ldr.4405 by Cochrane Thailand,

, Wiley Online Library on [06/08/2023]. See the Terms

and Condition:

(https:

onlinelibrary.wiley.com/term

-and-conditions) on Wiley Online Library for rules of use; OA

articles are governed by the applicable Creative Commons License

composting environments. When biochar is applied to the soil, its impact on soil physicochemical properties (e.g., porosity, water holding capacity, pH, and CEC) varies depending on biochar properties such as specific surface area, porosity, and functional groups (Sun et al., 2020). These changes caused by the different properties of biochar affect GHG emissions from the soil and other environmental processes.

Biochar feedstock is a key factor in determining biochar composition (Liu et al., 2019). In general, feedstock type affects the surface area, pH, and content of stable C in the biochars. For instance, owing to the higher content of lignin in wood biomass, biochar produced from wood typically has a higher surface area than that produced from grass and forms more organo-mineral layers to provide a nutrient shelter for microbes, thus improving microbial activities and changing soil GHG emissions (Hagemann et al., 2017). In contrast, biochars produced from feedstocks with higher cellulose and hemicellulose contents (e.g., sugarcane straw and rice husk) are characterized by higher pH values and nutrient concentrations (Higashikawa et al., 2016). For acidic paddy and wetland soils, the addition of alkaline biochar increases the soil pH (Sri et al., 2021). A higher soil pH is helpful for the growth of methanotrophs, resulting in decreased CH₄ emission from paddy soils (Dong et al., 2013). Applying biochar with high nutrient concentrations to the soil is conducive to increasing microbial nutrients and improving the activity of microorganisms. Moreover, a meta-analysis of 154 studies reported that biochars produced from biosolids had the best ability to retain nitrogen (N) in soils, followed by those produced from animal wastes. Compared with the biochars produced from animal wastes and biosolids, the woody and herbaceous biochars exhibited a better ability to mitigate N₂O emissions from soil (Li et al., 2019). There is abundant available N in animal waste and biosolid biochars, which may stimulate the growth of denitrifiers and contribute to N₂O emissions. Therefore, feedstock type should be considered an important factor affecting the properties of biochar when used for environmental applications and climate change mitigation. However, the results for biosolid-derived biochar are highly variable because of the diverse physicochemical properties of the feedstocks and the limited availability of studies on biosolidderived biochar. Therefore, further research on the impact of biosolidderived biochars on GHG emissions is needed to formulate comprehensive recommendations.

The pyrolysis temperature of biochar has been recognized as another important factor affecting its properties (Liu et al., 2019). As the pyrolysis temperature increases, the pH, electrical conductivity, ash content, and C stability of the biochar increase, whereas the yield of biochar decreases. Compared to biochars produced at medium (350°C-600°C) and high temperatures (>600°C), biochars produced at low temperatures (<350°C) generally contain a higher organic nutrient content which increases the co-metabolic interaction between biochars and microorganisms, thus resulting in the enhancement of microbial biomass and activities, especially for bacteria and fungi (Zhang et al., 2018). In addition, biochar produced at low temperatures (250-400°C) stimulates C mineralization, whereas biochar produced at high temperatures (525-650°C) suppresses C mineralization, ultimately decreasing CO_2 emissions (Wang, Wang, et al., 2019). However, high-temperature biochars may contain higher relative concentrations of toxic compounds (i.e., polycyclic aromatic hydrocarbons), affecting soil microbial biomass and activity (He et al., 2017). Simultaneously, the yield of high-temperature biochars was lower. Therefore, when choosing the biochar pyrolysis temperature, not only the impact of biochar on soil GHG emissions but also the cost savings of biochar production should be considered.

2.2 | Properties of engineered biochar relevant to climate change mitigation

Biochar properties can also be affected by post-treatment biochar production, that is, the production of engineered biochars. The properties of engineered biochars vary depending on the modification technologies, including physical (e.g., ball milling and magnetization) and chemical (e.g., acidification, alkalization, oxidation, and impregnation) methods (Panahi et al., 2020). Biochar modification is often used to increase its surface area, pore volume, surface functional groups, and surface chemistry properties. Through modification, biochar has a highly porous structure, which can improve a range of soil physical properties such as porosity and pore size distribution. This may further improve soil aeration, thereby stimulating the decomposition of soil organic C and the activity of methanotrophs (Liu et al., 2019). The engineering of biochar through ball milling has recently attracted significant research interest. Compared to pristine biochars, Nengineered biochar prepared by milling a mixture of biochar, bentonite, pregelatinized maize flour, and urea presents better environmental performance and lowers GHG emission intensity (Puga et al., 2020).

Few studies have reported the application of engineered biochars, including Fe-, N-, and phosphorus (P)-engineered biochars, in soil improvement. The biochar-supported FeS composite (FeS/biochar) can not only immobilize Cr(VI) through fractional precipitation in soil, but can also increase soil organic matter content, microbial activity, and CO₂ emissions (Lyu et al., 2018). As conductive and semiconductive materials, biochar and Fe may enhance direct interspecies electron transfer among soil microorganisms affecting GHG emissions (Liu et al., 2020). P-engineered biochars have improved stability owing to the formation of a P-containing compound that protects biochar C from oxidation (Guo & Chen, 2014). The co-pyrolysis of biomass with phosphate fertilizer could reduce C loss in soil. The role of minerals in biochar and their effects on biochar C stability are complex. Some inherent minerals in biochar can enhance the stability, whereas some extraneous minerals, such as Fe-bearing materials, reduce the stability of biochars. In contrast, inherent minerals can also reduce biochar C stability, whereas some extraneous minerals can enhance it (Buss et al., 2019). The incubation of biochar with soil minerals such as FeCl₃, AlCl₃, CaCl₂, and kaolinite could also increase the oxidation resistance of biochar (Yang et al., 2016). Clay types such as montmorillonite (MMT), red earth (RE), and bentonite have been used to synthesize engineered biochars as an efficient way to increase the stability of biochar in soil (Premarathna et al., 2019). Therefore, it is

³³³⁰ WILEY

TABLE 1 Overview of the main impact of biochar addition on upland soils

Biochar types	Soil type	Region	Influence	Reference
Sawdust biochar, <i>Sophora japonica</i> bark biochar	Obsidian and loess	Loess Plateau, China	 Soil CO₂ emission flux shows an increasing trend with an increase in the amount of biochar added Suppression of CH₄ emissions and no significant impact on N₂O emissions Differences exist among treatments with different biochars 	Guo et al. (2015)
Bamboo char	Dystric Cambisols	Obu City, Aichi Prefecture, Japan	• No significant changes in greenhouse gas (GHG) emissions	Watanabe et al. (2014)
Rice husks biochar	Orthic Anthrosols	Nanjing, China	 Increased CH₄ emissions Reduced NO₂ emissions Differences in GHG emissions between upland and wetland soils 	Wang et al. (2012)
Wheat straw biochar	Loamy soil	Central China Plain	 Reduced GHGs emissions Enhanced crop productivity	Zhang et al. (2012)
Barley straw biochar	Loamy soil	Sepung-ri, Gwangyang-eup, Gwangyang-si, and Jeollanam-do, South Korea	 Reduced N₂O emissions The combined treatment of biochar and chemical fertilizers was more effective in suppressing N₂O emissions than the treatment alone 	Kang et al. (2018)
Municipal solid waste biochar	/	Chongqing, China	 CO₂ emissions increased in the first 2 weeks Suppressed the total CO₂ emissions within 36 weeks 	Liu et al. (2015)
Ten types of biochar from Mediterranean agricultural residues	Sandy loam	Jumilla, Murcia, Spain	 CH₄ release was suppressed The starting material of biochar determined the difference in CH₄ release flux 	Pascual et al. (2020)
Fir sawdust	Luvisol soil	Luancheng, HeBei, China	 Suppressed the production of N₂O in the soil Stimulated the reduction of N₂O to N₂ 	Dong et al. (2020)
Wheat straw	Orthic Black Chernozem	Flagstaff County, southeast Alberta, Canada	 Reduced N₂O emissions No significant change in CO₂ or CH₄ emissions 	Wu et al. (2013)
Maize straw	Sandy loam soil	Fengqiu County, Henan Province, China	 Reduced N₂O emissions Reduced denitrification potential 	Niu et al. (2017)

important to develop new engineered biochars for better C sequestration and mitigation of GHG emissions. However, the relationship between stabilization and GHG emissions remains an interesting topic for further research.

3 | EFFECT OF BIOCHAR ON GHG EMISSIONS FROM VARIOUS ENVIRONMENTS

The addition of biochar could be used as a low-cost and highly efficient technology that might contribute to both climate change mitigation and adaptation (improving or maintaining soil quality), ensuring that the yield of upland and paddy crops is improved or maintained despite the changing climate (Pradhan et al., 2018). Reduced nitrogen loss, increased microbial activity, shorter time until maturity, and significantly less odour is observed when biochar is used as a compost amendment (Guo, Liu, & Zhang, 2020). As the physicochemical properties and microbial communities of upland soils, paddy and wetland soils, and compost are quite different, the application of biochar has different effects on GHG emissions in these three environments. Tables 1 and 2 summarize recent studies on GHG emissions resulting from the addition of biochar to upland fields, paddy fields, and wetland soils. In the following sections, we discuss how

-WILEY-

TABLE 2 O	verview of	the main i	mpact of	biochar add	lition on pa	ddv and wetland s	oils

Biochar type	Soil type	Region	Influence	Reference
Rice straw biochar	Rice paddy	Subtropical rice paddy China	 Biochar treatment decreased the cumulative CO₂ flux in the late paddy and for the complete year (early and late paddies) Biochar treatment also decreased the cumulative CH₄ flux in the early paddy 	Wang, Wang, et al. (2019)
Wheat straw biochar	Rice paddy	Dongshan Town, Suzhou City, Jiangsu Province, China	 The application of 4% biochar significantly increased N₂O emissions during the 45-day incubation by 291% and 256%, respectively The abundance and diversity of ammonia-oxidizing bacteria increased 	Lin et al. (2017)
Wheat straw biochar	Cadmium- and lead- contaminated rice paddy soil	Tai lake Plain, China	 No change in soil CO₂ emissions was observed at 10 t ha⁻¹ of biochar addition Biochar treatment reduced soil CO₂ emissions by 16%-24% at 20 and 40 t ha⁻¹ Biochar treatment increased rice yield by 25%-26% and thus enhanced ecosystem CO₂ sequestration by 47%-55% over the control Seasonal total N₂O emissions were reduced by 7.1%, 30.7%, and 48.6% under biochar addition at 10, 20, and 40 t ha⁻¹, respectively 	Zhang et al. (2015)
Mangrove biochar	Rice paddy	Rangbua, Chombueng District, Ratchaburi Province, Thailand	 Relative to control, biochar application reduced seasonal CH₄ emissions by 40.6% Biochar application enhanced soil organic carbon stock by 21.2% 	Sriphirom et al. (2020)
Rice straw biochar	Rice paddy	Yuhang District, Hangzhou, Zhejiang Province, China	• Biochar treatment reduced CH ₄ emissions under ambient conditions and significantly reduced emissions by 39.5% under simultaneously elevated temperature and CO ₂	Han et al. (2016)
Rice straw biochar	Rice paddy	Hwasungsi, Gyeonggido, Korea	 Biochar amendment did not significantly increase the CO₂ or CH₄ emissions Biochar addition increased the N₂O emissions The microbial biomass and the abundance of methane related microorganisms were not changed by biochar addition 	Yoo et al. (2015)
Wheat straw and sawdust biochars	Rice paddy	Taihu Lake region of China	 Biochar decreased CH₄ emissions Biochar application decreased N₂O emissions 	Zhou et al. (2018)
Rice chaff biochar	Rice paddy	Chunan-Si, Chungcheongnam-Do, Korea		Yoo et al. (2014)

3332 WILEY-

TABLE 2 (Continued)

Biochar type	Soil type	Region	Influence	Reference
			 Biochar treatment reduced the soil NH₄⁺ content and increased the NO₃⁻ content Biochar addition increased C contents in the wet stable aggregates of size 53-1000 mm, and the water holding capacity 	
Digested slurry biochar	Rice paddy	Graduate School of Horticulture, Chiba University, Matsudo, Japan	 Biochar treatment increased the CH₄ emissions Biochar treatment increased the soil NH₄⁺-N content 	Singla et al. (2014)
Tree branches biochar	Constructed wetlands	Yunnan University in Kunming, China	 Biochar treatment decreased the CO₂, CH₄, and N₂O emissions 	Ji, Chen, et al. (2020)
Cattail biochar (harvested from the wetland)	Constructed wetlands	Chongqing University, Chongqing, China	 Biochar reduced the global warming potential values of N₂O and CH₄ from 18.5% to 24.0% N₂O fluxes and global warming potential decreased, while CH₄ and CO₂ fluxes increased with increasing COD/N ratios 	Guo, Zhang, et al. (2020)
CO2 pli of biochar Acid Nataline Feedstock Wood waste Straw wast Straw wast T < 500°C		Biochar addition	Increase soil microbial biomass (more significant in upland soils) Increase oligotrophic bacteria Paddy and Paddy and	cow saturated ater content of upland soil improved but still dry) Decrease CO ₂ emissions



e

-40 -30 -20 -10 0 10 20 30 40 Change in GHGs intensity(%)

c(1)

-50

Paddy and

Composting

Pyrolysis < 400°C

450-500°C 550-600°C 600-800°C

Subtotal

Pyrolysis te T < 500°C T > 500°C

Feedstock Bark Chips Biochar C/N 30%-100% 100%-500%

Subtotal

Overall

perature

FIGURE 1 (a) Changes in CO₂ intensity in upland, rice paddy and wetlands, and composting environments after biochar application as influenced by pH, feedstock, pyrolysis temperature, application rate, and C/N ratio of biochar. The black solid line at zero indicates no change in CO₂ intensity after biochar addition. (b) the effects and mechanisms of biochar application on soil CO₂ emissions [Colour figure can be viewed at wileyonlinelibrary.com]

Decomposition

of unstable

carbon in

biochar

Adsorption

of CO₂

compost: Increase

copiotrophic bacteria

Increase

Decrease

(b)



FIGURE 2 (a) Changes in CH₄ intensity in upland, rice paddy and wetlands, and composting environments after biochar application as influenced by pH, feedstock, pyrolysis temperature, application rate, and C/N ratio of biochar. The black solid line at zero indicates no change in CH₄ intensity after biochar addition. (b) the effects and mechanisms of biochar application on soil CH₄ emissions [Colour figure can be viewed at wileyonlinelibrary.com]

biochar application affects the emissions of CO_2 , CH_4 , and N_2O in different environments. Moreover, a meta-analysis considering the interaction between these changes and GHG emissions is provided in Figures 1a–3a. Specifically, a literature search was conducted using Web of Science and Google Scholar databases from 1950 to 2021 using the keywords "biochar" AND "upland" OR "paddy" OR "composting" OR "greenhouse gas" OR "GHGs" OR "CO₂" OR "CH₄" OR "N₂O" OR "global warming potential (GWP)." Since most of the related studies separately assessed the effects of biochar application on GHGs emissions, physicochemical properties of biochar, and soil microbial properties, only 81 observations from 24 peer-reviewed studies were collected (listed in the supplemental material as Section 1). These results were discussed to clarify the different effects of biochar on GHGs emissions from upland, paddy, and wetland soils and composting environments.

Several parameters were identified as the main influencing factors in the meta-analysis, including pH of biochar, feedstock, pyrolysis temperature of biochar, application rate, C/N ratio of biochar, and experimental scale. An overall suppression effect among different environments was found, in the following order for different GHGs: $N_2O > CH_4 > CO_2$. Moreover, the addition of biochar can cause changes in soil physicochemical properties (bulk density, soil waterholding capacity, soil cation exchange capacity, pH, etc.), which affect soil microbial properties, including microbial biomass, microbial activity, and microbial community structure, which are related to GHG emissions in various environments (Guo, Liu, & Zhang, 2020). Herein, we summarize the microbial processes involved in the effects of biochar on GHGs emissions. The effects and mechanisms of biocharmediated GHG emissions are summarized in Figures 1b–3b.

3.1 | Effect of biochar on CO₂ emissions and its mechanism

As shown in Figure 1a, the meta-analysis results indicated that the overall reduction rate of CO_2 emissions intensity in the three different environmental processes of upland, rice paddies and wetlands, and composting was approximately 1%. However, the effect of biochar on the emission of CO_2 is quite different in the three different environments, showing an enhancing effect in uplands (a promotion rate of 9%, p < 0.05) and a suppression effect in paddy soil and composting processes (suppression rate of 10% and 2%, respectively, p < 0.05).

Specifically, as shown in Figure 1a, several parameters were identified in the meta-analysis as factors affecting CO₂ emissions in upland



FIGURE 3 (a) Changes in N₂O intensity in upland, rice paddy and wetlands, and composting environments after biochar application as influenced by pH, feedstock, pyrolysis temperature, application rate, and C/N ratio of biochar. The black solid line at zero indicates no change in N₂O intensity after biochar addition. (b) the effects and mechanisms of biochar application on soil N₂O emissions [Colour figure can be viewed at wileyonlinelibrary.com]

soil, including the pH of biochar, feedstock, pyrolysis temperature of biochar, application rate, and C/N ratio of biochar. Among them, the pH value and feedstocks showed a greater effect on increasing CO₂ emission intensity. That is, the addition of biochar promoted CO₂ emission intensity regardless of changes in pH and feedstocks of biochar (i.e., biochar pH and feedstock increased CO₂ emission intensity by 16% and 14%, respectively, p < 0.05). Moreover, the increase in soil CO₂ emission intensity was negatively correlated with biochar pyrolysis temperature, while positively correlated with biochar application rate and C/N. These results may be ascribed to the fact that lower pyrolysis temperature (500°C) results in more microbial available C and nutrients in biochar than a higher pyrolysis temperature (>500°C), which promotes high soil microbial activities to decompose soil organic matter and release more CO₂ from soil. At the same time, high temperature biochars (>500°C) may contain higher relative concentrations of toxic compounds (i.e., polycyclic aromatic hydrocarbons), which affect soil microbial biomass and activity (He et al., 2017). Overall, in upland soils, enzymes and labile organic matter are adsorbed from the bulk soil to the biochar surface, which is

more likely to cause significant microbial growth. In addition, the application of biochar to the soil directly affects the microbial community because of its unstable C components, which increase the apparent respiration rate of microorganisms and then increases soil CO_2 emissions (Irfan et al., 2019).

Unlike the trend in upland soils, the meta-analysis results showed that the addition of biochar usually decreases the cumulative CO_2 flux from paddy and wetland soils (Figure 1a). For example, compared to untreated paddy soils (a field experiment), the biochar-amended soils exhibited reduced CO_2 emissions (from 68,962 to 55,422 kg CO_2 -eq ha⁻¹) and increased rice yield (from 11.4 to 11.9 Mg ha⁻¹) (Wang, Wang, et al., 2019). Meta-analysis results suggested that biochar feed-stock, application rate, and pyrolysis temperature could influence CO_2 emissions from rice paddies and wetlands. For example, biochar from wood (a suppression rate of 35%, p < 0.05) can induce a greater suppression effect on CO_2 emissions than rice straw (a suppression rate of 12%, p < 0.05), probably because of the higher surface area and graphitic structure of biochar from wood (Hagemann et al., 2017), which is conducive to the suppression of soil organic carbon

mineralization and the adsorption of soil CO₂ molecules by biochar (Yu et al., 2021). The effect of the pyrolysis temperature of biochar on CO₂ emissions in rice paddies and wetlands is quite different. Compared with higher (600–800°C) and lower temperature (< 400°C) of biochars, which suppressed the CO₂ emissions intensity significantly, medium temperature biochars (450–600°C) had less suppression effect on CO₂ emissions (a suppression rate of 4% for 550–600°C, *p* < 0.05) and even greatly increased CO₂ emissions (a promotion rate of 45% for 450–500°C, *p* < 0.05). This may be due because medium pyrolysis temperature of biochars contain moderate organic nutrient content, pore structure, and surface area, and lower relative concentrations of toxic compounds, which increases the overall abundance and activities of microorganisms and promotes CO₂ emissions (Zhang et al., 2018).

The meta-analysis results showed that the addition of biochar to solid organic compost can regulate and mitigate CO2 emissions during composting (Figure 1a). The main influencing factors included pyrolysis temperature, raw materials, and initial C/N, all of which showed a low suppression effect on CO2 emissions (suppression rate of 0.1%-3.7%, p < 0.05). The result of this suppression comes from a combination of several reasons. For example, He et al. (2019) studied the effects of biochar on GHG emissions during composting in laboratory-scale composting systems, and found that the application of bamboo biochar reduced CO₂ emissions arising from composting (He et al., 2019). This has been ascribed to the biochar-mediated protection of organic matter against chemical oxidation and biological degradation (Ngo et al., 2013). Moreover, the addition of biochar to composting promotes enzyme activities (e.g., dehydrogenase, protease, cellulase, amylase, and xylanase) and reduces CO₂ emissions by affecting the carbon and nitrogen cycle (Awasthi et al., 2020). However, other studies have reported the opposite effects of biochar addition, that is, increased CO₂ emissions from the composting processes. The CO₂ emissions from chicken manure compost supplemented with biochar (27% w/w) increased by 6%-8% in smallscale laboratory composters (Chowdhury et al., 2014). This may be due to the high porosity and specific surface area of biochar, which allows a compost pile to have more oxygen to facilitate aeration, thus increasing CO₂ emissions (Wojciech et al., 2015). Other research indicated that higher CO₂ emissions during composting of mixtures amended with biochar could result from abiotic oxidation of biochar or biochar available carbon, which functions as an energy source for microorganisms (Dias et al., 2010).

Net ecosystem exchange of CO_2 (NEE) should also be considered when evaluating the effects of biochar amendment on soil CO_2 emissions. The NEE between terrestrial ecosystems and the atmosphere depends on the net C balance between the input and output of a given ecosystem and can be calculated as the difference between heterotrophic soil respiration and net primary production (Zhang et al., 2016). Azeem et al. (2019) conducted a 2-year field trial in an arid agricultural zone to investigate the effects of biochar on NEE for a legume-cereal crop rotation. The NEE for wheat decreased by 200% and 147% in the first year, and by 283% and 265% in the second year, and wheat yield increased by 6.2%–22.2% in soil amended with 0.25% and 0.5% biochar respectively (Azeem et al., 2019). The results revealed that biochar application improved the soil's physical and chemical properties, such as increasing the porosity and water-holding capacity of the soil (Major et al., 2010). As a result, biochar applications to soils enhanced crop productivity and limited nutrient leaching (Biederman & Harpole, 2013). However, no significant difference was observed for NEE in the first year of the mash bean crop; the NEE decreased by 46.8%–37.9% in the second season, and the mash bean yield increased by 3.9%–9.5%. The reason for this phenomenon may be that high rainfall during mash bean growing cycles leads to increased soil respiration, and the improvement of soil physical properties results in enhanced crop productivity which leads to no or small differences in NEE (Azeem et al., 2019).

The overall mechanism by which biochar regulates CO₂ emissions in various environments is illustrated in Figure 1b. Generally, the governing mechanisms, including both abiotic and biotic mechanisms, are summarized as follows: (1) the increase in soil pH and the high content of alkaline metals on the surface of biochar facilitates the precipitation of CO₂ to carbonates; (2) the adsorption of organic matter by biochar may be protected from further mineralization to produce CO_2 ; (3) the decrease in the abundance of two carbohydrate-mineralizing enzymes (glucosidase and cellobiosidase) reduces CO2 emissions; and (4) an increase in plant growth and plant biomass due to the addition of biochar increases the net exchange of CO_2 between the atmosphere and soil (Guo, Zhang, et al., 2020). Many researchers have demonstrated that soil pH is the main factor affecting the microbial community structure. Bacterial diversity was highest in neutral soils and lowest in acidic soils. Therefore, for paddy and wetland fields with lower pH, higher biochar addition led to a higher soil pH and bacterial diversity. For example, the enrichment of copiotrophic bacteria, such as Bacteroidetes and Gemmatimonadetes, and the decrease in oligotrophic bacteria, such as Acidobacteria in paddy and wetland fields were responsible for the decreased CO₂ emissions. However, when the biochar is added to the upland soil, the bacteria in the upland soil can adsorb to the surface of the biochar, making the bacteria in the soil less susceptible to soil leaching, thus increasing the number of bacteria in the soil. The biochar gaps are better able to protect microbes from competitors and thus enhance respiration of upland soil microbes in relation to soil available carbon (Li et al., 2021).

3.2 | Effect of biochar on CH₄ emissions and its mechanism

As shown in Figure 2a, the meta-analysis results confirmed that the addition of biochar generally suppressed the release of CH₄ from the three environments (upland soil, paddy and wetland fields, and compost) (Guo et al., 2015; Pascual et al., 2020), with an overall suppression of about 7% (p < 0.05). The suppression effect among different environments was in the following order: composting environment > rice paddies and wetlands > upland soil. It is speculated that the primary reason for this suppression is that the changes in the physical and chemical properties affect microbial activities. Biochar increases soil oxygen content because of its large pore structure. Since

methanogens are anaerobic bacteria, the aerated environment suppresses their activity, resulting in a decrease in the amount of CH₄ produced. However, the suppression of CH₄ emissions after biochar addition was not as strong in upland fields (a suppression rate of 3%, p < 0.05) as in other ecosystems (e.g., a suppression rate of 6% in rice paddies and wetlands), as shown in Figure 2a. This is because the low water content also suppresses the CH₄ oxidation process. Consequently, the promotion effect of biochar on methanotrophs in upland fields is weaker than in wet areas (Troy et al., 2013). Some studies have indicated that the addition of biochar can increase CH₄ emissions in uplands (Zhang et al., 2013). For example, a higher biochar application rate (>5 t ha⁻¹) provides a large amount of substrate, promoting the production of CH₄ (a promotion rate of 10%, p < 0.05), as confirmed by the results of the meta-analysis on CH₄ emission intensity in uplands (Figure 2a).

CH₄ emissions from rice paddies and wetlands were much higher than those from upland fields. Routine drainage and flooding of wetlands increase CH₄ emissions into the atmosphere. The meta-analysis results showed that the application of biochar to rice paddies and wetlands suppresses CH₄ emissions in general (Figure 2a). For example, in a 2-year field experiment conducted by Dong et al. (2013), rice straw and bamboo biochars were applied to paddy soils and CH₄ emissions were monitored for two growing seasons. The results showed that rice straw biochar had the most significant effect on the reduction of CH₄ emissions (causing 47.3%-86.4% reduction) and raised rice yield by 13.5%-6.1% during the two rice-growing cycles (Dong et al., 2013). This decrease may be ascribed to an increase in CH₄ oxidation and a decrease in methanogenic activity (Han et al., 2016). Specifically, biochar application decreases soil bulk density and increases soil aeration, thereby enhancing CH₄ oxidation (Liu et al., 2019). Moreover, soil pH is an important parameter to control soil CH₄ emission rates in paddy fields and wetlands because the biochemical activities of most methanogens are very sensitive to changes in soil pH. Soil pH increased after the addition of biochar. A higher soil pH was helpful for the growth of methanotrophs, resulting in reduced CH₄ emissions. However, different feedstocks of biochar have different effects on CH₄ emissions as shown in Figure 2a. Among them, biochar from straw suppressed CH₄ emissions intensity by 16% (p < 0.05), while biochar from wood significantly increased CH₄ emissions intensity by 34% (p < 0.05). The difference in the chemical properties of the biochars might explain this phenomenon. Compared with wood biochar, straw biochar generally has higher pH, which can significantly increase the degree of soil pH, increase the abundance of methane nutrient bacteria and promote methane oxidation (Dong et al., 2013). The higher pyrolysis temperature of biochar (>500°C) resulted in an enhanced inhibitory effect of biochar on CH₄ emissions. This was related to the soil redox potential, which also contributed to the reduction in CH_4 emissions. A soil redox potential of <-150 mV is beneficial to CH₄ production (Lyu et al., 2018). The addition of biochar might increase the redox potentials of paddy and wetland soils by affecting the water-holding capacity, soluble organic C, and metabolism of plant roots, thereby reducing CH₄ emissions. The biochar application rate and C/N ratio also had significant effects on CH₄

emissions in paddy and wetland soils. A higher application rate and lower C/N ratio are beneficial for suppressing CH_4 emissions which may be the result of both adsorption and microbial activity.

The meta-analysis results shown in Figure 2a confirmed that the addition of biochar significantly suppressed CH₄ emissions during composting (a suppression rate of 15%, p < 0.05) by improving the internal structure of compost piles, increasing the formation of aerobic sites, suppressing the activity of methanogens, enhancing the activity of methane-oxidizing bacteria, and reducing CH4 emissions (Sonoki et al., 2013). The proportion of biochar added was positively correlated with the reduction in CH₄ emissions during composting, for example, after 15 days of composting, the CH₄ emission concentration decreased from 1000 to 750 ppm as the biochar application rate increased from 5% to 20% (w/w) (Liu et al., 2017). Overall, the addition of biochar suppressed CH₄ emission intensity regardless of changes in application rate, feedstocks and C/N ratio of biochar, suggesting that biochar application is a good strategy to mitigate CH₄ emissions in composting systems. In addition to the reason previously mentioned that the addition of biochar improves the permeability of compost and changes the oxidation-reduction potential, so as to suppress the activity of methanogens and promote the activity of methane-oxidizing bacteria, another reason for biochar to reduce CH₄ emission in compost is its adsorption of $NH_{4}^{+}-N$, which decrease nitrogen availability to methanogens (Liu et al., 2017).

The overall mechanism by which biochar regulates CH₄ emission in various environments is shown in Figure 2b. Three processes determine the release of CH₄: CH₄ production, CH₄ oxidation, and CH₄ transport from soil to the atmosphere. CH_4 emissions are mainly related to the relative abundances of methanogens and methanotrophs, which are responsible for the production and oxidation of CH₄ respectively (Henri et al., 2018). The ratio of methanogens and methanotrophs is opposite to the suppression effect of biochar on CH₄ emission, which was confirmed in the paddy soil with long-time application of biochar (Wang, Shen, et al., 2019). In addition, two genes related to CH₄ emissions, mcrA (a methanogen) and pmoA (a methanotroph), have been well-studied, and there is a positive relationship between the copy number of mcrA and CH₄ emissions (Su et al., 2019). Methanogens are more active in weakly alkaline and neutral soils. Biochar addition generally results in an increase in pH and oxygen content in all three environments, which inhibited methanogens and reduced the emission of CH₄ (Pascual et al., 2020). Moreover, NO₃⁻-N was found to inhibit the activity of methanogens and enhance the activity of methanotroph (Nan et al., 2022). This indicates that the emission of CH₄ is related to N cycle by changing the relative abundance of different types of microbial community, which deserve to be studied further in the future.

3.3 | Effect of biochar on N₂O emissions and its mechanism

As shown in Figure 3a, the addition of biochar had the most obvious suppression effect on the release of N_2O from the three

environments (an overall suppression rate of 31%, p < 0.05) compared with CO₂ and CH₄ (Dong et al., 2020). The suppression effect on N₂O emission intensity among different environments was in the following order: upland soil (a suppression rate of 62%) > rice paddies and wetlands (a suppression rate of 20%) > composting environment (a suppression rate of 10%), which might be related to the different conditions. A meta-analysis of 208 peer-reviewed studies reported that biochar increased symbiotic biological N₂ fixation (63%), improved plant N uptake (11%), reduced soil N₂O emissions (32%), and decreased soil N leaching (26%) (Liu et al., 2018). However, the soil type was not considered in this meta-analysis.

In this work, the meta-analysis showed that N₂O emissions from upland were suppressed regardless of changes in pH of biochar, feedstock, pyrolysis temperature of biochar, application rate, and C/N ratio of biochar (Figure 3a). Alkaline conditions are favourable for N₂O emissions. In some cases, the pH value of upland soils is higher than that of biochar (Dong et al., 2020). When biochar is applied to upland fields, it decreases soil pH and reduces N₂O emissions. The pyrolysis temperature of biochar is also an important factor affecting N₂O emissions in upland soils. A 100-day laboratory incubation experiment by Pokharel et al. (2018) showed that a higher pyrolysis temperature of 550°C reduces N₂O emissions by 27.5%, while biochar pyrolyzes at 300°C without affecting N₂O emissions (Pokharel et al., 2018). These results are consistent with the meta-analysis results (Figure 3a) and might be ascribed to the fact that high-temperature biochar is more conducive to the transfer of electrons to soil-denitrifying microorganisms, leading to a more active N₂O reductase and an enhanced rate of N_2O reduction to N_2 . The application rate was positive, and the C/N ratio showed a negative suppression effect on N₂O emissions in upland soil, which can be related to the physicochemical properties of soil, such as soil porosity, pH value, and air permeability, which need to be comprehensively considered. For example, biochar application could enhance soil porosity to adsorb NH_4^+ while reducing NO_3^- produced by nitrification and N₂O produced by denitrification in soil (Zhang, Xiao, et al., 2020). Similarly, several studies have shown that biochar did not significantly reduce the release of N₂O and even promoted the release of N₂O in the absence of an external N source, indicating that N₂O release is regulated by soil saturated water content and plant N uptake (Zhang et al., 2012). Nevertheless, the application of biochar to upland fields is generally beneficial for alleviating the release of N₂O.

The reduction in soil N_2O emissions from paddy and wetland soils as a result of biochar application was confirmed in the meta-analysis

-WILEY <u>3337</u>

results, as shown in Figure 3a. The fluctuation of changes in N₂O emission intensity among different parameters, including pH of biochar, feedstock, pyrolysis temperature of biochar, application rate, and C/N ratio of biochar, was lower in paddy and wetland soils (20%, p < 0.05) than in upland soils (62%, p < 0.05). N₂O emissions are predominantly generated via N transformation in soils (Ji, Li, et al., 2020). The interaction between biochar and arbuscular mycorrhizal fungi affects N₂O emissions in paddy fields and constructed wetlands (Liang et al., 2019). This may be the reason why low pH favoured the suppression of N₂O emissions in paddy soils and constructed wetlands in the meta-analysis. The presence of arbuscular mycorrhizal fungi decreased the concentrations of chlorophyll and N in Phragmites australis and the concentrations of NH_4^+ -N, NO_3^- -N, inorganic N, and total N in paddy and wetland soils, thereby decreasing N₂O emissions. Moreover, biochar alters microbial activity and abundance, thereby affecting arbuscular mycorrhizal fungi and GHG emissions. In a field experiment, increasing the biochar application rate from 0 to 10 t ha^{-1} , 20 t ha^{-1} , and 40 t ha^{-1} reduced N₂O emissions from Cdand Pb-contaminated soils by 7.1%, 30.7%, and 48.6%, respectively, and increased rice yield by 10.0%, 25.1%, and 26.3% respectively (Zhang et al., 2015). The application rate of biochar was positively correlated with the suppression of N₂O emissions, which was consistent with the results of meta-analysis. Moreover, a meta-analysis of 88 studies on the effects of biochar on soil N2O emissions concluded that on average, biochar application resulted in a 38% reduction in N₂O emissions (Borchard et al., 2019).

The emissions of N₂O from composting can be reduced by applying biochar, although the changes in N₂O emission intensity are low (a suppression rate of 10%, p < 0.05) (Figure 3a). The availability of N in the compost is one of the main factors affecting N₂O generation. Biochar can limit N availability by directly adsorbing NO₃⁻ and NH₄⁺ and forming nutrient-rich organo-mineral complexes (Zwieten et al., 2010). Additionally, biochar can adsorb nitrate and dissolved organic C produced during composting, which can promote the complete denitrification of nitrate to produce N₂, thus reducing N₂O formation (Kammann et al., 2015). In contrast, N₂O emissions from composting during the maturation stage mainly depend on the degree of completion of the denitrification reactions and the proportion of biochar added (Wang et al., 2013). The addition of biochar increases the oxygen content in the compost pile and suppresses the activity of denitrifying bacteria, thereby weakening the completion of the denitrification reaction (Singh et al., 2010). In pilot-scale treatments (2 t of compost), Wang et al. (2013) found that an input of 3% (w/w) biochar

TABLE 3 Comparison of the mitigating effect of biochar in different environmental systems

System	CO ₂ emissions	N ₂ O emissions	CH ₄ emissions
Upland soils	Increase in the initial period, decrease in the long- term	Decrease	Small decrease or no effect
Rice paddy and wetlands	Changed based on conditions	Decrease	Decrease
Composting sites	Decrease	Decrease initially, increase at maturity	Decrease

reduces the abundance of denitrifying bacteria, suppresses denitrification, and reduces N₂O emissions. However, with a high biochar application rate (>8%), the compost accumulates a large amount of NH₄⁺-N, which facilitated the production of N₂O during the maturation stage.

The overall mechanism by which biochar regulates N₂O emission in various environments is summarized in Figure 3b. Generally, N₂O production involves two main microbial processes, nitrification and denitrification. N₂O-related functional genes, such as amoA, nirS, nirK, and nosZ, have been widely studied and can be used to elucidate ammonia oxidation, nitrification, and denitrification processes related to N₂O emissions (Harter et al., 2016). Variations in environmental factors, such as pH, oxygen, and water content, cause changes in the microbial activity involved in these two processes. There are five reasons for the reduction in N₂O release in soils as a result of biochar application. (1) Biochar reduces the activity of denitrifying bacteria and their enzymes by increasing soil porosity and permeability, thereby suppressing N₂O emissions (Singh et al., 2010). (2) Biochar may facilitate the transfer of electrons to soil-denitrifying microorganisms, leading to a more active N₂O reductase and an enhanced rate of N_2O reduction to N_2 (Cayuela et al., 2013). (3) Biochar produced from higher pyrolysis (>500°C) or feedstocks containing heavy metals may contain some toxic substances that inhibit nitrification and denitrification processes. (4) Biochar can immobilize the available N in the soil. thereby reducing the effective utilization of N in nitrification and denitrification reactions (Dong et al., 2020). (5) Furthermore, low soil pH is another reason for the reduction in N₂O emissions from soils. The increase in pH caused by the addition of biochar affects the activities of related enzymes involved in denitrification (such as nitrous oxide reductase), which is more pronounced in rice paddies, wetlands, and compositing systems (Cayuela et al., 2014).

4 | CONCLUSIONS AND FUTURE RESEARCH PROSPECTS

The widespread application of biochar and engineered biochar in different environments will affect soil and other biological processes that eventually affect GHG emissions. There has been much research on the application of biochar in the agricultural systems of upland and rice paddy fields and composting systems. Research to date suggests that biochar can improve soil quality by regulating soil pH, bulk density, water-holding capacity, and organic matter content, which can improve the agricultural productivity and quality of the composting process. Furthermore, biochar can regulate the emission of the three most important GHGs, that is, CO₂, N₂O, and CH₄ in different environmental processes. In conclusion, the effect of biochar on the emission of CO₂ in all three environments is still not clear, as contradictory results were obtained (Table 3). This can be attributed to the properties of the biochar itself and the negative and positive priming effects of biochar on the decomposition of soil organic matter. Despite this, it is becoming apparent that biochar can decrease the N₂O emissions from upland and paddy soils and compost environments to a certain

degree and considerably decrease the emission of CH₄ in paddy soils and composting systems. The emissions of CH₄ in upland soils did not appear to be affected by the addition of biochar. The emissions of CH₄ in upland soils increased because of the high pH value of some upland soils.

Based on our review, several future research directions are suggested:

- There are few reports on the effect of engineered biochar application on GHG emissions. Engineered biochars, such as nZVI-biochar composites, have been extensively studied and applied in the environment. The effects of these engineered biochars on GHG emissions differ from those of original biochars, and further studies are required. More importantly, engineered biochars can be developed with the primary or secondary aim of mitigating GHG emissions.
- A comprehensive analysis of the pyrolysis process, stability of biochar, and mitigation effect of biochar application should be carried out based on the concept of life cycle assessment (LCA) and compared with other processes, such as biogas fermentation, combustion, and direct application of biomass to the soil.
- 3. Owing to the large number of fertilizers, such as nitrogen and organic fertilizers used in agricultural production, it is possible to study the relationship between the change mechanism of soil microorganisms and the change of GHGs in the process of simultaneously applying fertilizer and biochar to different soils.
- 4. The effects of different feedstocks and production technologies of biochar on soil GHG emissions and crop yield need to be quantified and vetted. Guidelines on selecting and producing biochar formulations should be developed to improve soil health and environmental management, reduce the carbon footprint, abate climate change impacts on food production, and increase farmland profitability. Biochars can be tailored for specific applications through feedstock selection by modifying process conditions through pre- or post-production treatments to adjust pH by increasing nutrient levels and availability, carbon persistence, and adsorptive properties.

ACKNOWLEDGMENTS

This work was partially supported by: (1) the National Natural Science Foundation of China [42177218, 41877372, U18062162]; (2) the National Key R&D Program of China [2018YFC1802002]; (3) Hebei Outstanding Youth Science Foundation [D2019202453]; (4) China Postdoctoral Science Foundation [2019M660967]; and (5) the 111 Program, Ministry of Education, China [T2017002]. This work was also supported by the National Research Foundation of Korea (NRF) grant funded by the Korean Government (MSIT) (No. 2021R1A2C2011734), Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2021R1A6A1A10045235), and OJEong Resilience Institute, Korea University, Republic of Korea.

CONFLICT OF INTEREST

The author declares no competing financial interests.

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

ORCID

Jingchun Tang https://orcid.org/0000-0001-7633-1701 Scott X. Chang https://orcid.org/0000-0002-7624-439X Yong Sik Ok https://orcid.org/0000-0003-3401-0912

REFERENCES

- Aamer, M., Shaaban, M., Hassan, M. U., Huang, G., Liu, Y., & Tang, H. Y. (2020). Biochar mitigates the N₂O emissions from acidic soil by increasing the nosZ and nirK gene abundance and soil pH. *Journal of Environmental Management*, 255, 109891. https://doi.org/10.1016/j. jenvman.2019.109891
- Awasthi, M. K., Duan, Y., Awasthi, S. K., Liu, T., & Zhang, Z. (2020). Influence of bamboo biochar on mitigating greenhouse gas emissions and nitrogen loss during poultry manure composting. *Bioresource Technology*, 303, 122952. https://doi.org/10.1016/j.biortech.2020.122952
- Azeem, M., Hayat, R., Hussain, Q., Ahmed, M., Pan, G., Tahir, M. I., Imran, M., Irfan, M., & Mehmood-ul-Hassan. (2019). Biochar improves soil quality and N₂ fixation and reduces net ecosystem CO₂ exchange in a dryland legume-cereal cropping system. *Soil and Tillage Research*, 186, 172–182. https://doi.org/10.1016/j.still.2018.10.007
- Biederman, L. A., & Harpole, W. S. (2013). Biochar and its effects on plant productivity and nutrient cycling: A meta-analysis. GCB Bioenergy, 5(2), 202–214. https://doi.org/10.1111/gcbb.12037
- Borchard, N., Schirrmann, M., Luz Cayuela, M., Kammann, C., Wrage-Moennig, N., & Estavillo, J. M. (2019). Biochar, soil and land-use interactions that reduce nitrate leaching and N₂O emissions: A meta-analysis. Science of the Total Environment, 651(2), 2354–2364. https://doi. org/10.1016/j.scitotenv.2018.10.060
- Buss, W., Jansson, S., Wurzer, C., & Mašek, O. (2019). Synergies between BECCS and biochar–Maximizing carbon sequestration potential by recycling wood ash. ACS Sustainable Chemistry & Engineering, 7(4), 4204-4209. https://doi.org/10.1021/acssuschemeng.8b05871
- Cayuela, M. L., Sánchez-Monedero, M. A., Roig, A., Hanley, K., Enders, A., & Lehmann, J. (2013). Biochar and denitrification in soils: When, how much and why does biochar reduce N₂O emissions? *Scientific Reports*, 3, 1732. https://doi.org/10.1038/srep01732
- Cayuela, M. L., van Zwieten, L., Singh, B. P., Jeffery, S., Roig, A., & Sánchez-Monedero, M. A. (2014). Biochar's role in mitigating soil nitrous oxide emissions: A review and meta-analysis. Agriculture, Ecosystems & Environment, 191, 5–16. https://doi.org/10.1016/j.agee. 2013.10.009
- Chowdhury, M. A., de Neergaard, A., & Jensen, L. S. (2014). Potential of aeration flow rate and bio-char addition to reduce greenhouse gas and ammonia emissions during manure composting. *Chemosphere*, 97(2), 16–25. https://doi.org/10.1016/j.chemosphere.2013.10.030
- Dias, B. O., Silva, C. A., Higashikawa, F. S., Roig, A., & Sanchez-Monedero, M. A. (2010). Use of biochar as bulking agent for the composting of poultry manure: Effect on organic matter degradation and humification. *Bioresource Technology*, 101(4), 1239–1246. https://doi. org/10.1016/j.biortech.2009.09.024
- Dong, D., Yang, M., Wang, C., Wang, H., Li, Y., Luo, J., & Wu, W. (2013). Responses of methane emissions and rice yield to applications of biochar and straw in a paddy field. *Journal of Soils and Sediments*, 13(8), 1450–1460. https://doi.org/10.1007/s11368-013-0732-0
- Dong, W., Walkiewicz, A., Bieganowski, A., Oenema, O., Nosalewicz, M., He, C., Zhang, Y., & Hu, C. (2020). Biochar promotes the reduction of N₂O to N₂ and concurrently suppresses the production of N₂O in

calcareous soil. Geoderma, 362, 114091. https://doi.org/10.1016/j. geoderma.2019.114091

- Guo, F., Zhang, J., Yang, X., He, Q., Ao, L., & Chen, Y. (2020). Impact of biochar on greenhouse gas emissions from constructed wetlands under various influent chemical oxygen demand to nitrogen ratios. *Bioresource Technology*, 303, 122908. https://doi.org/10.1016/j.biortech. 2020.122908
- Guo, J., & Chen, B. (2014). Insights on the molecular mechanism for the recalcitrance of biochars: Interactive effects of carbon and silicon components. *Environmental Science & Technology*, 48(16), 9103–9112. https://doi.org/10.1021/es405647e
- Guo, X. X., Liu, H. T., & Zhang, J. (2020). The role of biochar in organic waste composting and soil improvement: A review. Waste Management, 102, 884–899. https://doi.org/10.1016/j.wasman.2019.12.003
- Guo, Y. L., Wang, D. D., Zheng, J. Y., Zhao, S. W., & Zhang, X. C. (2015). Effect of biochar on soil greenhouse gas emissions in semi-arid region. *Environmental Sciences*, 36(9), 3393–3400. https://doi.org/10.1016/j. chemosphere.2013.10.030
- Hagemann, N., Joseph, S., Schmidt, H. P., Kammann, C. I., Harter, J., Borch, T., Young, R. B., Varga, K., Taherymoosavi, S., & Elliott, K. W. (2017). Organic coating on biochar explains its nutrient retention and stimulation of soil fertility. *Nature Communications*, 8(1), 1089. https:// doi.org/10.1038/s41467-017-01123-0
- Han, X., Sun, X., Wang, C., Wu, M., Dong, D., Zhong, T., Thies, J. E., & Wu, W. (2016). Mitigating methane emission from paddy soil with rice-straw biochar amendment under projected climate change. *Scientific Reports*, 6, 24731. https://doi.org/10.1038/srep24731
- Harter, J., Weigold, P., El-Hadidi, M., Huson, D. H., Kappler, A., & Behrens, S. (2016). Soil biochar amendment shapes the composition of N₂O-reducing microbial communities. *Science of the Total Environment*, 562, 379–390. https://doi.org/10.1016/j.scitotenv.2016.03.220
- He, X., Yin, H., Han, L., Cui, R., Fang, C., & Huang, G. (2019). Effects of biochar size and type on gaseous emissions during pig manure/wheat straw aerobic composting: Insights into multivariate-microscale characterization and microbial mechanism. *Bioresource Technology*, 271, 375–382. https://doi.org/10.1016/j.biortech.2018.09.104
- He, Y., Zhou, X., Jiang, L., Li, M., Du, Z., Zhou, G., Shao, J., Wang, X., Xu, Z., Hosseini Bai, S., Wallace, H., & Xu, C. (2017). Effects of biochar application on soil greenhouse gas fluxes: A meta-analysis. *GCB Bioenergy*, 9(4), 743–755. https://doi.org/10.1111/gcbb.12376
- Henri, V. K., Bodelier, P., Ad Rian, H., Marion, M. F., & Veraart, A. J. (2018). Resistance and recovery of methane-oxidizing communities depends on stress regime and history; a microcosm study. *Frontiers in Microbiol*ogy, 9, 1714. https://doi.org/10.1038/s41467-017-01123-0
- Higashikawa, F. S., Conz, R. F., Colzato, M., Cerri, C. E. P., & Alleoni, L. R. F. (2016). Effects of feedstock type and slow pyrolysis temperature in the production of biochars on the removal of cadmium and nickel from water. *Journal of Cleaner Production*, 137, 965–972. https://doi.org/ 10.1016/j.jclepro.2016.07.205
- Huang, Y. F., Chiueh, P. T., Shih, C. H., Lo, S. L., Sun, L. P., Zhong, Y., & Qiu, C. S. (2015). Microwave pyrolysis of rice straw to produce biochar as an adsorbent for CO₂ capture. *Energy*, 84, 75–82. https://doi.org/ 10.1016/j.energy.2015.02.026
- Irfan, M., Hussain, Q., Khan, K. S., Akmal, M., Ijaz, S. S., Hayat, R., Khalid, A., Azeem, M., & Rashid, M. (2019). Response of soil microbial biomass and enzymatic activity to biochar amendment in the organic carbon deficient arid soil: A 2-year field study. Arabian Journal of Geosciences, 12(3), 95. https://doi.org/10.1007/s12517-019-4239-x
- Ji, B., Chen, J., Mei, J., Chang, J., Li, X., Jia, W., & Qu, Y. (2020). Roles of biochar media and oxygen supply strategies in treatment performance, greenhouse gas emissions, and bacterial community features of subsurface-flow constructed wetlands. *Bioresource Technology*, 302, 122890. https://doi.org/10.1016/j.biortech.2020.122890
- Ji, C., Li, S., Geng, Y., Miao, Y., Ding, Y., Liu, S., & Zou, J. (2020). Differential responses of soil N_2O to biochar depend on the predominant microbial

pathway. Applied Soil Ecology, 145, 103348. https://doi.org/10.1016/j. apsoil.2019.08.010

- Josep, B., Rafael, P., & Rodrigo, V. (2019). Automated measurements of greenhouse gases fluxes from tree stems and soils: Magnitudes, patterns and drivers. *Scientific Reports*, 9(1), 4005. https://doi.org/10. 1038/s41598-019-39663-8
- Kammann, C. I., Schmidt, H. P., Messerschmidt, N., Linsel, S., Steffens, D., Muller, C., Koyro, H. W., Conte, P., & Joseph, S. (2015). Plant growth improvement mediated by nitrate capture in co-composted biochar. *Scientific Reports*, 5, 11080. https://doi.org/10.1038/srep11080
- Kang, S. W., Kim, S. H., Park, J. H., Seo, D. C., Ok, Y. S., & Cho, J. S. (2018). Effect of biochar derived from barley straw on soil physicochemical properties, crop growth, and nitrous oxide emission in an upland field in South Korea. *Environmental Science and Pollution Research International*, 25(26), 25813–25821. https://doi.org/10.1007/s11356-018-1888-3
- Lee, J., Yang, X., Cho, S.-H., Kim, J.-K., Lee, S. S., Tsang, D. C. W., Ok, Y. S., & Kwon, E. E. (2017). Pyrolysis process of agricultural waste using CO₂ for waste management, energy recovery, and biochar fabrication. *Applied Energy*, 185, 214–222. https://doi.org/10.1016/j. apenergy.2016.10.092
- Li, C., Xiong, Y., Zou, J., Dong, L., Ren, P., & Huang, G. (2021). Impact of biochar and lignite-based amendments on microbial communities and greenhouse gas emissions from agricultural soil. *Vadose Zone Journal*, 20(2), 20105. https://doi.org/10.1002/vzj2.20105
- Li, S. M., Harris, S., Anandhi, A., & Chen, G. (2019). Predicting biochar properties and functions based on feedstock and pyrolysis temperature: A review and data syntheses. *Journal of Cleaner Production*, 215, 890– 902. https://doi.org/10.1016/j.jclepro.2019.01.106
- Li, Y. F., Hu, S. D., Chen, J. H., Muller, K., Li, Y. C., Fu, W. J., Lin, Z. W., & Wang, H. L. (2018). Effects of biochar application in forest ecosystems on soil properties and greenhouse gas emissions: A review. *Journal of Soils and Sediments*, 18(2), 546–563. https://doi.org/10.1007/s11368-017-1906-y
- Lian, F., & Xing, B. (2017). Black carbon (biochar) in water/soil environments: Molecular structure, sorption, stability, and potential risk. Environmental Science & Technology, 51(23), 13517–13532. https://doi. org/10.1021/acs.est.7b02528
- Liang, J. F., An, J., Gao, J. Q., Zhang, X. Y., Song, M. H., & Yu, F. H. (2019). Interactive effects of biochar and AMF on plant growth and greenhouse gas emissions from wetland microcosms. *Geoderma*, 346, 11– 17. https://doi.org/10.1016/j.geoderma.2019.03.033
- Lin, Y. X., Ding, W. X., Liu, D. Y., He, T. H., Yoo, G., Yuan, J. J., Chen, Z. M., & Fan, J. L. (2017). Wheat straw-derived biochar amendment stimulated N₂O emissions from rice paddy soils by regulating the amoA genes of ammonia-oxidizing bacteria. *Soil Biology and Biochemistry*, 113, 89–98. https://doi.org/10.1016/j.soilbio.2017.06.001
- Liu, G., Xie, M., & Zhang, S. (2015). Effect of organic fraction of municipal solid waste (OFMSW)-based biochar on organic carbon mineralization in a dry land soil. *Journal of Material Cycles and Waste Management*, 19(1), 473–482. https://doi.org/10.1007/s10163-015-0447-y
- Liu, N., Zhou, J., Han, L., Ma, S., Sun, X., & Huang, G. (2017). Role and multi-scale characterization of bamboo biochar during poultry manure aerobic composting. *Bioresource Technology*, 241, 190–199. https:// doi.org/10.1016/j.biortech.2017.03.144
- Liu, Q., Zhang, Y., Liu, B., Amonette, J. E., Lin, Z., Liu, G., Ambus, P., & Xie, Z. (2018). How does biochar influence soil N cycle? A meta-analysis. *Plant and Soil*, 426(1–2), 211–225. https://doi.org/10.1007/ s11104-018-3619-4
- Liu, T., Wang, Y., Liu, C., Li, X., Cheng, K., Wu, Y., Fang, L., Li, F., & Liu, C. (2020). Conduction band of hematite can mediate cytochrome reduction by Fe(II) under dark and anoxic conditions. *Environmental Science & Technology*, 54(8), 4810–4819. https://doi.org/10.1021/acs.est. 9b06141

- Liu, X., Mao, P., Li, L., & Ma, J. (2019). Impact of biochar application on yield-scaled greenhouse gas intensity: A meta-analysis. *Science of the Total Environment*, 656, 969–976. https://doi.org/10.1016/j.scitotenv. 2018.11.396
- Lyu, H., Zhang, Q., & Shen, B. (2020). Application of biochar and its composites in catalysis. *Chemosphere*, 240, 124842. https://doi.org/10. 1016/j.chemosphere.2019.124842
- Lyu, H., Zhao, H., Tang, J., Gong, Y., Huang, Y., Wu, Q., & Gao, B. (2018). Immobilization of hexavalent chromium in contaminated soils using biochar supported nanoscale iron sulfide composite. *Chemosphere*, 194, 360–369. https://doi.org/10.1016/j.chemosphere.2017.11.182
- Major, J., Lehmann, J., Rondon, M., & Goodale, C. (2010). Fate of soilapplied black carbon: Downward migration, leaching and soil respiration. *Global Change Biology*, 16(4), 1366–1379. https://doi.org/10. 1111/j.1365-2486.2009.02044.x
- Nan, Q., Fang, C., Cheng, L., Hao, W., & Wu, W. (2022). Elevation of NO₃⁻⁻N from biochar amendment facilitates mitigating paddy CH₄ emission stably over seven years. *Environmental Pollution*, 295, 118707. https://doi.org/10.1016/j.envpol.2021.118707
- Ngo, P. T., Rumpel, C., Ngo, Q. A., Alexis, M., Velasquez Vargas, G., Mora Gil, M. L., Dang, D. K., & Jouquet, P. (2013). Biological and chemical reactivity and phosphorus forms of buffalo manure compost, vermicompost and their mixture with biochar. *Bioresource Technology*, 148, 401–407. https://doi.org/10.1016/j.biortech.2013.08.098
- Niu, Y., Chen, Z., Müller, C., Zaman, M. M., Kim, D., Yu, H., & Ding, W. (2017). Yield-scaled N₂O emissions were effectively reduced by biochar amendment of sandy loam soil under maize-wheat rotation in the North China Plain. Atmospheric Environment, 170, 58–70. https://doi. org/10.1016/j.atmosenv.2017.09.050
- Panahi, H. K. S., Dehhaghi, M., Ok, Y. S., Nizami, A.-S., Khoshnevisan, B., Mussatto, S. I., Aghbashlo, M., Tabatabaei, M., & Lam, S. S. (2020). A comprehensive review of engineered biochar: Production, characteristics, and environmental applications. *Journal of Cleaner Production*, 270, 122462. https://doi.org/10.1016/j.jclepro.2020.122462
- Pariyar, P., Kumari, K., Jain, M. K., & Jadhao, P. S. (2020). Evaluation of change in biochar properties derived from different feedstock and pyrolysis temperature for environmental and agricultural application. *Science of the Total Environment*, 713, 136433. https://doi.org/10. 1016/j.scitotenv.2019.136433
- Pascual, M. B., Sanchez-Monedero, M. A., Chacon, F. J., Sanchez-Garcia, M., & Cayuela, M. L. (2020). Linking biochars properties to their capacity to modify aerobic CH₄ oxidation in an upland agricultural soil. *Geoderma*, 363, 114179. https://doi.org/10.1016/j.geoderma.2020. 114179
- Pokharel, P., Kwak, J. H., Yong, S. O., & Chang, S. X. (2018). Pine sawdust biochar reduces GHG emission by decreasing microbial and enzyme activities in forest and grassland soils in a laboratory experiment. *Science of the Total Environment*, 625, 1247–1256. https://doi.org/10. 1016/j.scitotenv.2017.12.343
- Pradhan, A., Chan, C., Roul, P. K., Halbrendt, J., & Sipes, B. (2018). Potential of conservation agriculture (CA) for climate change adaptation and food security under rainfed uplands of India: A transdisciplinary approach. Agricultural Systems, 163, 27–35. https://doi.org/10.1016/j. agsy.2017.01.002
- Premarathna, K. S. D., Rajapaksha, A. U., Adassoriya, N., Sarkar, B., Sirimuthu, N. M. S., Cooray, A., Ok, Y. S., & Vithanage, M. (2019). Claybiochar composites for sorptive removal of tetracycline antibiotic in aqueous media. *Journal of Environmental Management*, 238, 315–322. https://doi.org/10.1016/j.jenvman.2019.02.069
- Puga, A. P., Grutzmacher, P., Cerri, C. E. P., Ribeirinho, V. S., & Andrade, C. A. (2020). Biochar-based nitrogen fertilizers: Greenhouse gas emissions, use efficiency, and maize yield in tropical soils. *Science* of the Total Environment, 704, 135375. https://doi.org/10.1016/j. scitotenv.2019.135375

- Singh, B. P., Hatton, B. J., Singh, B., Cowie, A. L., & Kathuria, A. (2010). Influence of biochars on nitrous oxide emission and nitrogen leaching from two contrasting soils. *Journal of Environmental Quality*, 39(4), 1224–1235. https://doi.org/10.2134/jeq2009.0138
- Singla, A., Dubey, S. K., Singh, A., & Inubushi, K. (2014). Effect of biogas digested slurry-based biochar on methane flux and methanogenic archaeal diversity in paddy soil. Agriculture, Ecosystems & Environment, 197, 278–287. https://doi.org/10.1016/j.agee.2014.08.010
- Song, C., Liu, Q., Deng, S., Li, H., & Kitamura, Y. (2019). Cryogenic-based CO₂ capture technologies: State-of-the-art developments and current challenges. *Renewable and Sustainable Energy Reviews*, 101, 265–278. https://doi.org/10.1016/j.rser.2018.11.018
- Sonoki, T., Furukawa, T., Jindo, K., Suto, K., Aoyama, M., & Sanchez-Monedero, M. A. (2013). Influence of biochar addition on methane metabolism during thermophilic phase of composting. *Journal of Basic Microbiology*, 53(7), 617–621. https://doi.org/10.1002/jobm. 201200096
- Spokas, K. A., & Reicosky, D. C. (2009). Impacts of sixteen different biochars on soil greenhouse gas production. Annals of Environmental Science, 3(612), 179-193. https://www.researchgate.net/publication/48856254_Impacts_ of_sixteen_different_biochars_on_soil_greenhouse_gas_production
- Sri, S. S., Palanivelu, K., Ramachandran, A., & Raghavan, V. (2021). Biochar from biomass waste as a renewable carbon material for climate change mitigation in reducing greenhouse gas emissions-a review. *Biomass Conversion and Biorefinery*, 11, 2247–2267. https://doi.org/10.1007/ s13399-020-00604-5
- Sriphirom, P., Chidthaisong, A., Yagi, K., Tripetchkul, S., & Towprayoon, S. (2020). Evaluation of biochar applications combined with alternate wetting and drying (AWD) water management in rice field as a methane mitigation option for farmers' adoption. *Soil Science & Plant Nutrition*, 66(1), 235–246. https://doi.org/10.1080/00380768.2019. 1706431
- Su, X., Wang, Y., He, Q., Hu, X., & Chen, Y. (2019). Biochar remediates denitrification process and N₂O emission in pesticide chlorothalonilpolluted soil: Role of electron transport chain. *Chemical Engineering Journal*, 370, 587–594. https://doi.org/10.1016/j.cej.2019.03.195
- Sun, H., Feng, Y., Xue, L., Mandal, S., Wang, H., Shi, W., & Yang, L. (2020). Responses of ammonia volatilization from rice paddy soil to application of wood vinegar alone or combined with biochar. *Chemosphere*, 242, 125247. https://doi.org/10.1016/j.chemosphere. 2019.125247
- Thangarajan, R., Bolan, N. S., Kunhikrishnan, A., Wijesekara, H., Xu, Y., Tsang, D. C. W., Song, H., Ok, Y. S., & Hou, D. (2018). The potential value of biochar in the mitigation of gaseous emission of nitrogen. *Sci*ence of the Total Environment, 612, 257–268. https://doi.org/10.1016/ j.scitotenv.2017.08.242
- Troy, S. M., Lawlor, P. G., O'Flynn, C. J., & Healy, M. G. (2013). Impact of biochar addition to soil on greenhouse gas emissions following pig manure application. *Soil Biology and Biochemistry*, 60, 173–181. https://doi.org/10.1016/j.soilbio.2013.01.019
- Wang, C., Lu, H., Dong, D., Deng, H., Strong, P. J., Wang, H., & Wu, W. (2013). Insight into the effects of biochar on manure composting: Evidence supporting the relationship between N₂O emission and denitrifying community. *Environmental Science & Technology*, 47(13), 7341– 7349. https://doi.org/10.1021/es305293h
- Wang, C., Shen, J. L., Liu, J. Y., Qin, H. L., Yuan, Q., Fan, F. L., Hu, Y. J., Wang, J., Wei, W. X., Li, Y., & Wu, J. S. (2019). Microbial mechanisms in the reduction of CH4 emission from double rice cropping system amended by biochar: A four-year study. *Soil Biology and Biochemistry*, 135, 251–263. https://doi.org/10.1016/j.soilbio.2019.05.012
- Wang, C., Wang, W., Sardans, J., Singla, A., Zeng, C., Lai, D. Y. F., & Penuelas, J. (2019). Effects of steel slag and biochar amendments on CO₂, CH₄, and N₂O flux, and rice productivity in a subtropical Chinese paddy field. *Environmental Geochemistry and Health*, 41(3), 1419– 1431. https://doi.org/10.1007/s10653-018-0224-7

- Wang, J., Pan, X., Liu, Y., Zhang, X., & Xiong, Z. (2012). Effects of biochar amendment in two soils on greenhouse gas emissions and crop production. *Plant and Soil*, 360(1–2), 287–298. https://doi.org/10.1007/ s11104-012-1250-3
- Watanabe, A., Ikeya, K., Kanazaki, N., Makabe, S., Sugiura, Y., & Shibata, A. (2014). Five crop seasons records of greenhouse gas fluxes from upland fields with repetitive applications of biochar and cattle manure. *Journal of Environmental Management*, 144, 168–175. https://doi.org/ 10.1016/j.jenvman.2014.05.032
- Wojciech, C., Krystyna, M., Rafaela, C., Janczak, D., & Andrzej, L. (2015). Co-composting of poultry manure mixtures amended with biochar-the effect of biochar on temperature and C-CO₂ emission. *Bioresource Technology*, 200, 921. https://doi.org/10.1016/j.biortech.2015.11.019
- Woolf, D., Amonette, J. E., Street-Perrott, F. A., Lehmann, J., & Joseph, S. (2010). Sustainable biochar to mitigate global climate change. *Nature Communications*, 1, 56. https://doi.org/10.1038/ncomms1053
- Wu, F., Jia, Z., Wang, S., Chang, S. X., & Startsev, A. (2013). Contrasting effects of wheat straw and its biochar on greenhouse gas emissions and enzyme activities in a Chernozemic soil. *Biology and Fertility of Soils*, 49(5), 555–565. https://doi.org/10.1007/s00374-012-0745-7
- Xiao, Y., Lyu, H. H., Tang, J. C., Wang, K., & Sun, H. W. (2020). Effects of ball milling on the photochemistry of biochar: Enrofloxacin degradation and possible mechanisms. *Chemical Engineering Journal*, 384, 123311. https://doi.org/10.1016/j.cej.2019.123311
- Yang, F., Zhao, L., Gao, B., Xu, X., & Cao, X. (2016). The interfacial behavior between biochar and soil minerals and its effect on biochar stability. *Environmental Science & Technology*, 50(5), 2264–2271. https://doi. org/10.1021/acs.est.5b03656
- Yoo, G., Kim, H., Chen, J., & Kim, Y. (2014). Effects of biochar addition on nitrogen leaching and soil structure following fertilizer application to rice paddy soil. Soil Science Society of America Journal, 78(3), 852–860. https://doi.org/10.2136/sssaj2013.05.0160
- Yoo, G., Kim, Y. J., Lee, Y. O., & Ding, W. (2015). Investigation of greenhouse gas emissions from the soil amended with rice straw biochar. *KSCE Journal of Civil Engineering*, 20(6), 2197–2207. https://doi.org/ 10.1007/s12205-015-0449-2
- Yu, H., Wang, T., Huang, Q., Song, K., Zhang, G., Ma, J., & Xu, H. (2021). Effects of elevated CO₂ concentration on CH₄ and N₂O emissions from paddy fields: A meta-analysis. *Science China Earth Sciences*, 65(1), 96–106. https://doi.org/10.1007/s11430-021-9848-2
- Zhang, A., Bian, R., Hussain, Q., Li, L., Pan, G., Zheng, J., Zhang, X., & Zheng, J. (2013). Change in net global warming potential of a ricewheat cropping system with biochar soil amendment in a rice paddy from China. Agriculture, Ecosystems and Environment, 173, 37–45. https://doi.org/10.1016/j.agee.2013.04.001
- Zhang, A., Bian, R., Li, L., Wang, X., Zhao, Y., Hussain, Q., & Pan, G. (2015). Enhanced rice production but greatly reduced carbon emission following biochar amendment in a metal-polluted rice paddy. *Environmental Science and Pollution Research International*, 22(23), 18977–18986. https://doi.org/10.1007/s11356-015-4967-8
- Zhang, A. F., Liu, Y. M., Pan, G. X., Hussain, Q., Li, L. Q., Zheng, J. W., & Zhang, X. H. (2012). Effect of biochar amendment on maize yield and greenhouse gas emissions from a soil organic carbon poor calcareous loamy soil from Central China plain. *Plant and Soil*, 351(1–2), 263–275. https://doi.org/10.1007/s11104-011-0957-x
- Zhang, L., Jing, Y., Xiang, Y., Zhang, R., & Lu, H. (2018). Responses of soil microbial community structure changes and activities to biochar addition: A meta-analysis. Science of the Total Environment, 643, 926–935. https://doi.org/10.1016/j.scitotenv.2018.06.231
- Zhang, M., & Ok, Y. S. (2014). Biochar soil amendment for sustainable agriculture with carbon and contaminant sequestration. *Carbon Management*, 5(3), 255–257. https://doi.org/10.1080/17583004.2014. 973684
- Zhang, Q., Xiao, J., Xue, J., & Zhang, L. (2020). Quantifying the effects of biochar application on greenhouse gas emissions from agricultural

3342 WILEY-

papermill waste on agronomic performance and soil fertility. *Plant and Soil*, 327(1), 235–246. https://doi.org/10.1007/s11104-009-0050-x

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Lyu, H., Zhang, H., Chu, M., Zhang, C., Tang, J., Chang, S. X., Mašek, O., & Ok, Y. S. (2022). Biochar affects greenhouse gas emissions in various environments: A critical review. *Land Degradation & Development*, 33(17), 3327–3342. https://doi.org/10.1002/ldr.4405

soils: A global meta-analysis. Sustainability, 12(8), 3436. https://doi.org/10.3390/su12083436
 Zhang, X., Qu, J. S., Li, H., La, S. K., Tian, Y. Q., & Gao, L. H. (2020). Biochar

- addition combined with daily fertigation improves overall soil quality and enhances water-fertilizer productivity of cucumber in alkaline soils of a semi-arid region. *Geoderma*, *363*, 114170. https://doi.org/10. 1016/j.geoderma.2019.114170
- Zhang, Y., Lin, F., Wang, X., Zou, J., & Liu, S. (2016). Annual accounting of net greenhouse gas balance response to biochar addition in a coastal saline bioenergy cropping system in China. *Soil and Tillage Research*, 158, 39–48. https://doi.org/10.1016/j.still.2015.11.006
- Zhou, B., Feng, Y., Wang, Y., Yang, L., Xue, L., & Xing, B. (2018). Impact of hydrochar on rice paddy CH₄ and N₂O emissions: A comparative study with pyrochar. *Chemosphere*, 204, 474–482. https://doi.org/10.1016/ j.chemosphere.2018.04.056
- Zwieten, L. V., Kimber, S., Morris, S., Chan, K. Y., Downie, A., Rust, J., Joseph, S., & Cowie, A. (2010). Effects of biochar from slow pyrolysis of

and Conditions

: (https://onlinelibrary.wiley.com/term)

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

1099145x, 2022, 17, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/dr.4405 by Cochrane Thailand, Wiley Online Library on [06/08/2023]. See the Terms